Original Article

Clin Shoulder Elbow 2020;23(2):94-99
<https://doi.org/10.5397/cise.2020.00052>

eISSN 2288-8721

Comparison of open reduction and internal fixation with total elbow arthroplasty for intra-articular distal humeral fractures in older age: a retrospective study

Jong Seok Baik, Sung Hyun Lee, Hyun Tak Kang, Tae Hyun Song, Jeong Woo Kim

Department of Orthopedic Surgery, Wonkwang University Hospital, Wonkwang University School of Medicine, Iksan, Korea

Background: Intra-articular distal humeral fractures can be surgically challenging. It remains under discussion whether open reduction and internal fixation (ORIF) or total elbow arthroplasty (TEA) is more beneficial for treatment of the elderly. This study aimed to compare the clinical and functional outcomes of ORIF and TEA for managing intra-articular distal humerus fractures in patients aged 65 years or older.

Methods: Patients who underwent ORIF (n=28) or TEA (n=43) for intra-articular distal humerus fracture between May 2008 and December 2018 were reviewed. Range of motion, Mayo Elbow Performance Score (MEPS), Disabilities of the Arm, Shoulder, and Hand (DASH) score, radiologic outcomes, and surgical complications were evaluated at the final follow-up visit.

Results: The ORIF and TEA groups showed a mean arc of flexion-extension of $97^{\circ} \pm 21^{\circ}$ and $101^{\circ} \pm 12^{\circ}$, respectively. The mean MEPS and DASH scores were 94 ± 15 and 27 ± 12 points, respectively, in the ORIF group and 81 ± 27 and 47 ± 28 points in the TEA group. This difference was statistically significant. The incidence of total complications was similar between the groups.

Conclusions: In patients older than 65 years with intra-articular distal humerus fracture, ORIF had better outcomes than TEA.

Keywords: Fracture fixation; Total elbow replacement; Humeral fracture; Aged, 65 and over; Treatment outcome

INTRODUCTION

Distal humeral fractures account for 2% of all fractures in adults [1]. In younger patients, these fractures are mainly caused by high-energy trauma, whereas, among older patients, they are more often the result of a direct injury that occurs from falling. Anatomical joint reconstruction and stable fixation of intra-articular and comminuted fractures remain difficult to achieve, and the operation for displaced intra-articular distal humeral fractures is partic-

ularly challenging.

Open reduction and internal fixation (ORIF) with double-locking plate osteosynthesis has become the gold standard for fixation of intra-articular distal humeral fractures, though the optimal positioning and configuration of the plates remain controversial [2-4]. In addition, elbow stiffness, malunion, nonunion, fixation failure, and ulnar neuropathy are common sequelae, with an overall complication rate greater than 35% [5-9]. In older individuals with low bone densities, there is an increased risk of osteosynthesis failure [10].

Received: January 31, 2020   Revised: March 22, 2020   Accepted: March 22, 2020

Correspondence to: Jeong Woo Kim

Department of Orthopedic Surgery, Wonkwang University School of Medicine, 460 Iksan-daero, Iksan 54538, Korea

Tel: +82-63-859-1360, Fax: +82-63-852-9329, E-mail: serina@wonkwang.ac.kr, ORCID: <https://orcid.org/0000-0002-0828-7179>

IRB approval: Wonkwang University Hospital (No. WKUHIRB 2018-01-011)

Financial support: None.

Conflict of interest: None.

Copyright© 2020 Korean Shoulder and Elbow Society. All Rights Reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Total elbow arthroplasty (TEA) is considered effective for treating nonunion, malunion, posttraumatic arthritis, and posttraumatic instability, with the literature reporting good to excellent results in 85% to 90% of patients at 5 to 10 years after surgery [11-13]. However, TEA for these injuries is technically difficult, and the complication rates are higher than those for elbows that have not undergone prior operations [7,11]. The purpose of this study was to compare the clinical and functional outcomes of ORIF with those of TEA for displaced intra-articular distal humeral fractures in patients aged 65 years and older.

METHODS

This study was approved by Institutional Review Board of Wonkwang University Hospital (IRB No. WKUHIRB 2018-01-011), and it was exempted from informed consent. We retrospectively reviewed the medical records of patients who underwent ORIF or TEA for intra-articular distal humeral fractures between May 2008 and December 2018. Eligible patients were those that met the following inclusion criteria: (1) aged of 65 years or older; (2) displaced and comminuted intra-articular distal humeral fractures requiring surgical treatment (AO/Orthopaedic Trauma Association [OTA] classification type 13C) [14], (3) surgical treatment with either ORIF with double-locking plate osteosynthesis or TEA, and (4) a minimum follow-up of 2 years.

Patients with extra- or partial-articular distal humeral fractures (AO/OTA classification types 13A and 13B), intra-articular distal humeral fractures not requiring surgical treatment (AO/OTA classification type 13C), open fractures, vascular injuries, previous ipsilateral distal humeral fractures, pathologic fractures, or fractures with diaphyseal extension ≥ 8 cm were excluded. A total of 77 patients met the inclusion criteria of this study. Six patients died from

unrelated causes. Of the remaining 71 patients, 28 underwent ORIF with double-locking plate osteosynthesis, while 43 underwent TEA. Patient demographics are summarized in Table 1.

Patient medical records were reviewed for demographic and perioperative data. All patients were invited for a clinical follow-up examination comprising evaluation of range of motion, elbow stability, and neurologic deficits. Additionally, pain status, patient satisfaction, and revision surgeries were documented. Finally, the Mayo Elbow Performance Score (MEPS) and Disabilities of the Arm, Shoulder, and Hand (DASH) score were examined. The functional outcome, based on the arc of motion, MEPS, and DASH score, as well as minor and major complications were evaluated and compared between the TEA and ORIF groups. All follow-up assessments were conducted by the same surgeon who performed the surgery.

Surgical Techniques

All surgeries were performed by the same senior surgeon (JWK).

ORIF Group

ORIF was performed using a posterior approach. After exposing the triceps muscle, the elbow joint was revealed by chevron osteotomy of the olecranon for accurate anatomical reduction. After temporary K-wire fixation of the fracture fragments, double-locking plates were applied to the medial and lateral sides of the distal humerus. The ulnar nerve was transposed into an anterior subcutaneous position at the conclusion of all surgical procedures. After postoperative immobilization for two days, passive range-of-motion exercise was performed for 6 weeks.

TEA Group

A Coonrad-Morrey semiconstrained prosthesis (Zimmer Biomet, Warsaw, IN, USA) was used in all cases. A midline triceps split or

Table 1. Patient demographics

Variable	ORIF	TEA	P-value
Sex (male:female)	7 (25):21 (75)	9 (21):34 (79)	0.59
Age (yr)	76 (65–89)	79 (65–91)	0.16
BMI (kg/m ²)	24 $\pm$ 3 (19–30)	23 $\pm$ 4 (15–34)	0.49
DM	4 (14)	8 (18.6)	0.34
Smoking	2 (7)	2 (4.6)	0.28
Follow-up period (mo)	31 (9–58)	34 (6–116)	0.59
AO classification			0.31
C1	8	10	
C2	2	10	
C3	18	23	

Values are presented as number (%), mean (range), or mean $\pm$ standard deviation (range).

ORIF: open reduction and internal fixation; TEA: total elbow arthroplasty; BMI: body mass index; DM: diabetes mellitus.

triceps-sparing approach was used. With the triceps-sparing approach, the surgeon used the working space created by condylar resection to perform TEA without detaching the triceps from the olecranon [11,15]. The ulnar nerve was transposed into an anterior subcutaneous position at the conclusion of all surgical procedures. After postoperative immobilization for 2 days, early functional mobilization was started without weight-bearing for 6 weeks.

Statistical Analysis

Statistical analyses were performed using IBM SPSS ver. 22.0 (IBM Corp., Armonk, NY, USA). The functional outcomes were statistically compared using the Mann-Whitney U-test as a two-way analysis of variance for independent factors.

RESULTS

Clinical Outcomes

In the ORIF group, the mean pain score was 3.1 (range, 1–6), mean MEPS was 94 (range, 75–100), and mean DASH score was 27 (range, 10–45). Based on MEPS, there were 20 excellent and seven good results. With respect to motion, average extension was 17° (range, 5°–40°), average flexion was 120° (range, 80°–140°), and average arc of flexion–extension was 97° (range, 70°–130°). Mean supination angle was 84° (range, 80°–90°), and mean pronation angle was 88° (range, 75°–90°) (Table 2).

All radiographs of fractures that healed without revision (all patients) showed anatomical reduction with stable fixation (Fig. 1). We did not intentionally rule out fractured patients who needed revision, but there was no patient who required it.

Table 2. Functional results comparing ORIF and TEA

Variable	ORIF	TEA	P-value
Pain (NRS)	3.1 (1–6)	3.5 (0–8)	0.21
MEPS	94 ± 15 (75–100)	81 ± 27 (50–85)	0.028*
DASH score	27 ± 12 (10–45)	47 ± 28 (10–75)	0.038*
ROM (°)			
Extension	17 ± 21 (5–40)	15 ± 3 (0–30)	0.104
Flexion	120 ± 18 (100–140)	124 ± 15 (80–150)	0.265
Flexion–extension	97 ± 21 (70–130)	101 ± 12 (80–140)	0.089
Supination	84 ± 6 (80–90)	84 ± 6 (80–90)	0.126
Pronation	88 ± 5 (75–90)	89 ± 6 (80–90)	0.072
Satisfaction			-
Excellent	20 (71.4)	5 (11.6)	
Good	7 (25)	18 (41.8)	
Fair	1 (3.5)	14 (32.6)	
Poor	0	6 (14.0)	

Values are presented as mean (range), mean ± standard deviation (range), or number (%).

ORIF: open reduction and internal fixation, TEA: total elbow arthroplasty, NRS: numeric rating scale, MEPS: Mayo Elbow Performance Score, DASH: Disabilities of the Arm, Shoulder, and Hand, ROM: range of motion.

*Statistically significant.

Fig. 1. Serial X-rays of open reduction and internal fixation (ORIF) case. (A) Anteroposterior and lateral X-rays of a 66-year-old female who suffered an AO classification 13C distal humerus fracture with a fall. (B) An X-ray performed after ORIF, showing dual-plate fixation. (C) Bone union after 6 months.

Patients with infection were treated with antibiotics and did not need additional surgery. One patient had Brooker type III heterotopic ossification in the brachialis musculature.

In the TEA group, mean pain score was 3.5 (range, 0–8), mean MEPS was 81 (range, 50–85), and mean DASH score was 47 (range, 10–75). Based on MEPS, there were five excellent and 18 good results. With respect to motion, average extension was 15° (range, 0°–30°), average flexion was 124° (range, 80°–150°), and average arc of flexion–extension was 101° (range, 80°–140°). Mean supination angle was 84° (range, 80°–90°), and mean pronation angle was 89° (range, 80°–90°) (Table 2).

At the final follow-up, incomplete radiolucent lines were seen around the humeral implant in six cases and in the ulnar implant in five cases. Wear of the polyethylene bushings at the hinge was directly correlated with follow-up duration. When we calculated satisfaction, the ORIF group (96.4%) contained a higher percentage of patients who answered good or excellent compared with the TEA group (53.4%).

Complications

Complications occurred in 13 patients in the ORIF group and 14 patients in the TEA group (Table 3). Two patients underwent revision in the TEA group, including one due to deep infection and one due to periprosthetic fracture with aseptic loosening of the humeral stem. These two cases in the TEA group were separated from the other, nonrevised TEA cases when measuring the clinical scores. The single case of deep infection occurred seven years after the initial TEA in a female patient. She underwent implant removal, insertion of antibiotic beads, and an installation of new prosthesis with antibiotic-loaded acrylic cement in the first-stage surgery. Six weeks after the first-stage surgery, revision TEA was performed after confirming no infection. Separately, two cases of superficial infection occurred in the ORIF group and were treated with intra-

venous antibiotics without surgery.

In the TEA group, the case of periprosthetic fracture occurred around the tip of the humeral implant, which had completely loosened. Here, the humeral implant was replaced with a longer one and augmented using a humerus shaft strut allograft (Fig. 2). Another case of loosening occurred between the cement and both the humerus and the ulna, while screw-loosening occurred in one case in the ORIF group. Also, neurological complications occurred in nine cases, including two in the TEA group and seven in the ORIF group. These patients reported only dysesthesia of the fourth and fifth fingers without sensory or motor deficits, with all recovering. Heterotopic ossification occurred in eight patients, but there was no instability or other clinical symptoms, and no surgical treatment was required.

We classified aseptic loosening, fractures, and infection as major complications; thus, the ORIF group had mainly minor complications such as ulnar nerve symptoms. When we calculated the rate of major complications of total complications, three of 13 (23.1%) in the ORIF group and four of 14 (28.6%) in the TEA group had major complications. Thus, the rate was higher in the TEA group, although there was no statistical significance.

DISCUSSION

The purpose of this study was to compare the clinical and func-

Table 3. Complications

Variable	ORIF	TEA	P-value
Wound dehiscence	0	1 (2.3)	
Aseptic loosening	1 (3.5)	2 (4.7)	
Fracture	0	1 (2.3)	
Heterotrophic ossification	1 (3.5)	7 (16.3)	
Ulnar nerve symptom	7 (25)	2 (4.7)	
Infection	2 (7.1)	1 (2.3)	
Elbow stiffness*	2 (7.1)	0	
Total	13 (46)	14 (32)	0.18

Values are presented as number (%).

ORIF: open reduction and internal fixation, TEA: total elbow arthroplasty.

*Flexion <120° and loss of extension >30°.

Fig. 2. A 75-year-old female suffered AO classification 13C distal humerus fracture with a slip down (A, B) and was treated with total elbow arthroplasty (TEA) (C, D). After 3 years, the patient complained of upper arm pain after carrying a heavy load. Periprosthetic fracture with aseptic loosening around the humeral stem was observed (E, F). Revision TEA with longer humeral stem and humerus shaft strut allograft was performed (G, H). The final functional outcome was relatively good.

tional outcomes of ORIF with those of TEA for displaced intra-articular distal humerus fractures in patients aged 65 years or older. At the time of final follow-up, the ORIF group had significantly higher MEPS and DASH scores compared with the TEA group.

Although the recommended treatment for comminuted and displaced intra-articular distal humerus fractures is osteosynthesis, this method can be technically difficult, especially in older patients [16,17]. Previous data on ORIF for distal humerus fractures indicate that complications such as fixation failure, persistent pain and/or stiffness, heterotopic ossification, ulnar nerve entrapment, non-union, malunion, and posttraumatic arthritis are common in all age groups.

Several studies have suggested that primary TEA is a reliable treatment for severe intra-articular distal humerus fractures in elderly patients. Morrey et al. [18] reported a series of 21 patients with a mean age of 72 years who underwent primary TEA for comminuted distal humerus fractures. They reported good or excellent results in 95% of the patients at a mean follow-up of 3.3 years, with a reoperation rate of 5% (one elbow). Gambirasio et al. [19] reported 10 women (mean age, 85 years) who underwent primary TEA. Of them, eight had an excellent outcome and two had a good outcome based on MEPS. Garcia et al. [20] evaluated 16 patients with a mean age of 73 years (range, 61–95 years) who underwent primary TEA with a mean follow-up of 3 years (range, 1.5–5 years). These authors reported mean DASH score of 23 (range, 1–63) and mean MEPS of 93 (range, 80–100). Frankle et al. [21] reported 11 excellent, one good, zero fair, and zero poor results in patients who underwent TEA.

There are several studies comparing ORIF and TEA for comminuted intra-articular distal humerus fractures. Frankle et al. [21] reported better functional outcomes and higher MEPS in the TEA group as well as a higher incidence of complications in the ORIF group. In a prospective multicenter study, McKee et al. [22] reported improvement in MEPS and DASH scores in the TEA group relative to in the ORIF group. Egol et al. [23] reported no significant differences between the groups in a study comparing TEA and locking-plate osteosynthesis. Obrebsky et al. [24] reported no strong evidence to choose one of two treatments for deployment in patients in their 60s who have a long predicted remaining lifespan and no comorbidities.

Ulnar neuropathy is a common complication after surgery. McKee et al. [22] reported that postoperative ulnar neuropathy was the single most common complication in their study comparing TEA and ORIF for distal humerus fractures. In our study, ulnar neuropathy was also the most common complication (nine cases) and was significantly more frequently observed in the ORIF group. We note that these complications occurred despite routinely trans-

posing the ulnar nerve in all patients. Thus, the surgeon should be extremely cautious of the ulnar nerve during surgery. Rates of major complications, such as infection and loosening, were similar in the two groups.

Infection seems to occur due to swelling; soft tissue damage after trauma; or shear force during early range of motion or due to the large volume of the dual plate, which can irritate the skin because the elbow joint has thin soft tissue. In our study, the infection rate of the ORIF group (7.1%) was not as high as reported in other researches [25]. Proper soft tissue control during the operation should be ensured.

Limitations of our study, including its retrospective design and the small number of patients, are comparable to those of previous studies but might have underpowered our study. Furthermore, despite a mean follow-up length of 34 months, it is difficult to compare the preoperative status between groups because only the final follow-up results were evaluated.

In older patients with intra-articular distal humeral fractures, those undergoing ORIF had better clinical outcomes and similar complication rates than those receiving TEA. As life expectancy increases, many TEA patients become too old to undergo surgery when the lifespan of their prostheses ends, resulting in many cases where patients have no choice but to bear the worn-out prosthesis for longer, i.e., up to 20 to 30 years. Therefore, rather than performing primary TEA right away just because the patient is elderly, if it is possible to perform osteosynthesis, ORIF can be a better choice for older patients with intra-articular distal humeral fractures.

ORCID

Jong Seok Baik	https://orcid.org/0000-0002-1877-1837
Sung Hyun Lee	https://orcid.org/0000-0001-6468-1456
Hyun Tak Kang	https://orcid.org/0000-0001-8050-3246
Tae Hyun Seong	https://orcid.org/0000-0001-9629-4856
Jeong Woo Kim	https://orcid.org/0000-0002-0828-7179

REFERENCES

1. Robinson CM, Hill RM, Jacobs N, Dall G, Court-Brown CM. Adult distal humeral metaphyseal fractures: epidemiology and results of treatment. *J Orthop Trauma* 2003;17:38-47.
2. Zalavras CG, Vercillo MT, Jun BJ, Otarodifard K, Itamura JM, Lee TQ. Biomechanical evaluation of parallel versus orthogonal plate fixation of intra-articular distal humerus fractures. *J Shoulder Elbow Surg* 2011;20:12-20.
3. Shin SJ, Sohn HS, Do NH. A clinical comparison of two different

- double plating methods for intraarticular distal humerus fractures. *J Shoulder Elbow Surg* 2010;19:2-9.
4. Hackl M, Wegmann K, Bartsch V, Ries C, Burkhart KJ, Muller LP. Impingement of the posterolateral humeral plate: landmarks for prevention in double-plate osteosynthesis of distal humeral fractures. *Obere Extremitat* 2014;9:186-91.
  5. Korner J, Lill H, Muller LP, et al. Distal humerus fractures in elderly patients: results after open reduction and internal fixation. *Osteoporos Int* 2005;16:S73-9.
  6. McKee MD, Wilson TL, Winston L, Schemitsch EH, Richards RR. Functional outcome following surgical treatment of intra-articular distal humeral fractures through a posterior approach. *J Bone Joint Surg Am* 2000;82:1701-7.
  7. McKee MD, Jupiter JB. Fractures of the distal humerus. In: Browner B, Jupiter J, Levine A, Trafton P, eds. *Skeletal trauma*. 3rd ed. Philadelphia, PA: Lippincott; 2002. p. 765-82.
  8. O'Driscoll SW. Optimizing stability in distal humeral fracture fixation. *J Shoulder Elbow Surg* 2005;14 Suppl S:186S-194S.
  9. Sodergard J, Sandelin J, Bostman O. Postoperative complications of distal humeral fractures. 27/96 adults followed up for 6 (2-10) years. *Acta Orthop Scand* 1992;63:85-9.
  10. Korner J, Lill H, Muller LP, Rommens PM, Schneider E, Linke B. The LCP-concept in the operative treatment of distal humerus fractures: biological, biomechanical and surgical aspects. *Injury* 2003;34 Suppl 2:B20-30.
  11. McKee MD, Pugh DM, Richards RR, Pedersen E, Jones C, Schemitsch EH. Effect of humeral condylar resection on strength and functional outcome after semiconstrained total elbow arthroplasty. *J Bone Joint Surg Am* 2003; 85:802-7.
  12. Morrey BF, Adams RA, Bryan RS. Total replacement for post-traumatic arthritis of the elbow. *J Bone Joint Surg Br* 1991; 73:607-12.
  13. Morrey BF, Adams RA. Semiconstrained elbow replacement for distal humeral nonunion. *J Bone Joint Surg Br* 1995;77:67-72.
  14. Fracture and dislocation compendium: Orthopaedic Trauma Association Committee for Coding and Classification. *J Orthop Trauma* 1996;10 Suppl 1:v-ix. 1-154.
  15. Armstrong AD, Yamaguchi K. Total elbow arthroplasty and distal humerus elbow fractures. *Hand Clin* 2004;20:475-83.
  16. John H, Rosso R, Neff U, Bodoky A, Regazzoni P, Harder F. Operative treatment of distal humeral fractures in the elderly. *J Bone Joint Surg Br* 1994;76:793-6.
  17. Jupiter JB, Neff U, Holzach P, Allgower M. Intercondylar fractures of the humerus. An operative approach. *J Bone Joint Surg Am* 1985;67:226-39.
  18. Morrey BF, Chao E, Cooney W, et al. Indications in patient selection. In: Morrey BF, ed. *Joint replacement arthroplasty*. New York, NY: Churchill Livingstone; 1991. p. 275-91.
  19. Gambirasio R, Riand N, Stern R, Hoffmeyer P. Total elbow arthroplasty as primary treatment for distal humeral fractures in elderly patients. *J Bone Joint Surg Br* 2001; 83:974-78.
  20. Garcia JA, Mykula R, Stanley D. Complex fractures of the distal humerus in the elderly: the role of total elbow replacement as primary treatment. *J Bone Joint Surg Br* 2002;84:812-6.
  21. Frankle MA, Herscovici D Jr, DiPasquale TG, Vasey MB, Sanders RW. A comparison of open reduction and internal fixation and primary total elbow arthroplasty in the treatment of intraarticular distal humerus fractures in women older than age 65. *J Orthop Trauma* 2003;17:473-80.
  22. McKee MD, Veillette CJ, Hall JA, et al. A multicenter, prospective, randomized, controlled trial of open reduction--internal fixation versus total elbow arthroplasty for displaced intra-articular distal humeral fractures in elderly patients. *J Shoulder Elbow Surg* 2009;18:3-12.
  23. Egol KA, Tsai P, Vazques O, Tejwani NC. Comparison of functional outcomes of total elbow arthroplasty vs plate fixation for distal humerus fractures in osteoporotic elbows. *Am J Orthop (Belle Mead NJ)* 2011;40:67-71.
  24. Obrebsky WT, Bhandari M, Dirschl DR, Schemitsch E. Internal fixation versus arthroplasty of comminuted fractures of the distal humerus. *J Orthop Trauma* 2003;17:463-5.
  25. Savvidou OD, Zampeli F, Koutsouradis P, et al. Complications of open reduction and internal fixation of distal humerus fractures. *EFORT Open Rev* 2018;3:558-67.